

KLOE RESULTS ON LIGHT MESON PROPERTIES

FEDERICO NGUYEN for the KLOE COLLABORATION^a
 INFN “Roma TRE”, Roma (Italy) – e-mail address: nguyen@fis.uniroma3.it

The KLOE experiment operating at the ϕ -factory DAΦNE has collected an integrated luminosity of about 2.5 fb^{-1} and 250 pb^{-1} , on and off the ϕ meson peak respectively. Recent results achieved from studying properties of pseudoscalar and scalar mesons are presented.

1 Introduction: KLOE detector

The KLOE detector consists of a cylindrical drift chamber¹ (3.3 m length and 2 m radius), surrounded by a calorimeter² made of lead and scintillating fibers. The detector is inserted in a superconducting coil producing a magnetic field $B=0.52 \text{ T}$. Large angle tracks from the origin ($\theta > 45^\circ$) are reconstructed with momentum resolution $\sigma_p/p = 0.4\%$. Photon energies and times are measured with resolutions of $\sigma_E/E = 5.7\%/\sqrt{E(\text{GeV})}$ and $\sigma_t = 57 \text{ ps}/\sqrt{E(\text{GeV})} \oplus 100 \text{ ps}$.

2 Pseudoscalar Mesons

KLOE collected a statistics of about 10^8 η and 5×10^5 η' events, produced through magnetic dipole transitions, $\phi \rightarrow \eta\gamma$ and $\phi \rightarrow \eta'\gamma$, with full reconstruction of decay products.

$\eta \rightarrow \pi^+\pi^-e^+e^-$. CP violating (CPV) mechanisms can be tested in the decays of the pseudoscalar mesons by measuring an asymmetry $\mathcal{A}_\phi$ in the angle ϕ between the e^+e^- and $\pi^+\pi^-$ planes in the meson rest frame. In the η case, a nonzero $\mathcal{A}_\phi$ value would signal CPV dynamics not directly related to most widely studied flavour changing neutral current processes. Furthermore, possible contributions^{3,4} beyond the Standard Model can raise $\mathcal{A}_\phi$ up to $\mathcal{O}(10^{-2})$.

The $\eta \rightarrow \pi^+\pi^-e^+e^-$ analysis⁵ is based on a sample of 1.7 fb^{-1} . The event selection consists of the requirement of one photon of $E > 250 \text{ MeV}$ energy, namely the monochromatic photon from $\phi \rightarrow \eta\gamma$, and four charged tracks coming from the interaction region. Mass assignment for each track is done using time of flight of the charged particles measured in the calorimeter. Background sources are due to ϕ decays: $\phi \rightarrow \pi^+\pi^-\pi^0$ or $\eta \rightarrow \pi^+\pi^-\pi^0$ processes with π^0 Dalitz decay; or *continuum processes*: $e^+e^- \rightarrow e^+e^-\gamma$ events with γ conversions, studied using data taken at $\sqrt{s} = 1 \text{ GeV}$. The contamination is evaluated by fitting the sidebands of the $M_{\pi\pi ee}$ data

^aF. Ambrosino, A. Antonelli, M. Antonelli, F. Archilli, C. Bacci, P. Beltrame, G. Bencivenni, S. Bertolucci, C. Bini, C. Bloise, S. Bocchetta, F. Bossi, P. Branchini, P. Campana, G. Capon, T. Capussela, F. Ceradini, S. Chi, G. Chiefari, P. Ciambrone, F. Crucianelli, E. De Lucia, A. De Santis, P. De Simone, G. De Zorzi, A. Denig, A. Di Domenico, C. Di Donato, B. Di Micco, A. Doria, M. Dreucci, G. Felici, A. Ferrari, M. L. Ferrer, S. Fiore, C. Forti, P. Franzini, C. Gatti, P. Gauzzi, S. Giovannella, E. Gorini, E. Graziani, W. Kluge, V. Kulikov, F. Lacava, G. Lanfranchi, J. Lee-Franzini, D. Leone, M. Martini, P. Massarotti, W. Mei, S. Meola, S. Miscetti, M. Moulson, S. Müller, F. Murtas, M. Napolitano, F. Nguyen, M. Palutan, E. Pasqualucci, A. Passeri, V. Patera, F. Perfetto, M. Primavera, P. Santangelo, G. Saracino, B. Sciascia, A. Sciubba, A. Sibidanov, T. Spadaro, M. Testa, L. Tortora, P. Valente, G. Venzoni, R. Versaci, G. Xu.

spectrum with background components after loose cuts on the kinematic fit χ^2 and on the sum of momenta of the charged particles. Signal events are computed after rejecting γ conversions, and from the fit the branching ratio is evaluated: $\text{BR}(\eta \rightarrow \pi^+\pi^-e^+e^-) = (26.8 \pm 0.9_{\text{stat}} \pm 0.7_{\text{sys}}) \times 10^{-5}$. The asymmetry $\mathcal{A}_\phi$ is measured from momenta of the charged particles and background




Figure 1: Left: $M_{\pi\pi ee}$ distributions around the η mass resulting from the fit: $\chi^2/\text{dof} = 32.5/30$. Middle: $\sin\phi \cos\phi$ distributions for $535 < M_{\pi\pi ee} < 555$ MeV. Right: $Z_{\eta'}^2$ vs. φ_P , with constraints from $Y1 = \Gamma(\eta' \rightarrow \gamma\gamma)/\Gamma(\pi^0 \rightarrow \gamma\gamma)$, $Y2 = \Gamma(\eta' \rightarrow \rho\gamma)/\Gamma(\omega \rightarrow \pi^0\gamma)$, $Y3 = R_\phi$ and $Y4 = \Gamma(\eta' \rightarrow \omega\gamma)/\Gamma(\omega \rightarrow \pi^0\gamma)$ measurements.

subtracted. From the selected sample of 1555 events, the first measurement of $\mathcal{A}_\phi$ performed is found compatible with zero at the 3% level: $\mathcal{A}_\phi = (-0.6 \pm 2.5_{\text{stat}} \pm 1.8_{\text{sys}}) \times 10^{-2}$. Left and middle panels of Fig. 1 show the $M_{\pi\pi ee}$ spectrum after the fit and the $\sin\phi \cos\phi$ distribution in the signal region.

η - η' mixing. From the selection of $\phi \rightarrow \eta'\gamma \rightarrow \pi^+\pi^-\gamma$ and $\phi \rightarrow \eta\gamma \rightarrow \gamma\gamma$ events out of a sample of 430 pb^{-1} , the ratio $R_\phi \equiv \Gamma(\phi \rightarrow \eta'\gamma)/\Gamma(\phi \rightarrow \eta\gamma)$ is measured. η , η' are known to be linear combinations of strange, $s\bar{s}$, and nonstrange, $n\bar{n}$, quarks. Moreover the η' could contain a gluonium component GG , so that

$$\begin{aligned} |\eta'\rangle &= X_{\eta'}|n\bar{n}\rangle + Y_{\eta'}|s\bar{s}\rangle + Z_{\eta'}|GG\rangle & X_{\eta'} &= \cos\varphi_G \sin\varphi_P \\ |\eta\rangle &= \cos\varphi_P|n\bar{n}\rangle - \sin\varphi_P|s\bar{s}\rangle & Y_{\eta'} &= \cos\varphi_G \cos\varphi_P \\ & & Z_{\eta'} &= \sin\varphi_G \end{aligned}$$

where φ_P is the pseudoscalar mixing angle. A combined fit⁶ of the KLOE measured value of $R_\phi \propto \cot^2\varphi_P \cos^2\varphi_G$ with available measurements of $\Gamma(\eta' \rightarrow \rho\gamma)/\Gamma(\omega \rightarrow \pi^0\gamma)$, $\Gamma(\eta' \rightarrow \gamma\gamma)/\Gamma(\pi^0 \rightarrow \gamma\gamma)$ and $\Gamma(\eta' \rightarrow \omega\gamma)/\Gamma(\omega \rightarrow \pi^0\gamma)$, yields the following results, if $Z_{\eta'}^2$ is fixed or not:

$$\varphi_P = (41.5^{+0.6}_{-0.7})^\circ, \quad Z_{\eta'}^2 = 0 \text{ fixed}, \quad P_{\chi^2} = 0.01, \quad (1)$$

$$\varphi_P = (39.7 \pm 0.7)^\circ, \quad Z_{\eta'}^2 = 0.14 \pm 0.04, \quad P_{\chi^2} = 0.49. \quad (2)$$

A 3σ evidence is found for gluonium content in η' , as shown in Fig. 1 right panel. Further analyses, including the new measurement⁷ of the $\text{BR}(\omega \rightarrow \pi^0\gamma)$ done by KLOE, confirm the gluonium content in the η' wavefunction.

3 Scalar Mesons

The still unresolved structure of these states is studied either through electric dipole transitions such as $\phi \rightarrow a_0(980)\gamma$ and looking at the mass spectrum of the scalar meson decay products, or with the search for processes like $\phi \rightarrow [a_0(980) + f_0(980)]\gamma \rightarrow K\bar{K}\gamma$ and $\gamma\gamma \rightarrow \sigma(600) \rightarrow \pi\pi$.

$\phi \rightarrow a_0(980)\gamma \rightarrow \eta\pi^0\gamma$. Two independent analyses⁸ using $\eta \rightarrow 2\gamma$ or $\eta \rightarrow \pi^+\pi^-\pi^0$ decays are performed from a sample of 410 pb⁻¹. Both analyses share the requirement of five photons from the interaction point. The selection of also two tracks of opposite charge, while less efficient for $\eta \rightarrow \pi^+\pi^-\pi^0$ events, has a selected sample with smaller background than from the $\eta \rightarrow 2\gamma$ channel. The absence of a major source of interfering background allows to obtain the branching fraction directly from event counting. Table 1 shows that the two samples lead to consistent branching ratio values, thus a combined fit of the two spectra is performed. The couplings, fitted according to the Kaon Loop⁹ and the No Structure¹⁰ models, point to a total width in the range $[80 \div 105]$ MeV and to a sizeable $s\bar{s}$ content of the $a_0(980)$.

channel features	$\eta \rightarrow \gamma\gamma$	$\eta \rightarrow \pi^+\pi^-\pi^0$
signal efficiency	40%	20%
$B/(S+B)$	50%	14%
$\text{BR}(\phi \rightarrow \eta\pi^0\gamma) \times 10^5$	7.01(10)(20)	7.12(13)(22)
fit parameter	Kaon Loop	No Structure
m_{a_0} (MeV)	982.5(1.6)(1.1)	982.5 (fixed)
$g_{a_0 K^+ K^-}$ (GeV)	2.15(6)(6)	2.01(7)(28)
$g_{a_0 \eta \pi}$ (GeV)	2.82(3)(4)	2.46(8)(11)
$g_{\phi a_0 \gamma}$ (GeV ⁻¹)	–	1.83(3)(8)
χ^2/dof	157.1/136	140.6/133
CL	10.4%	30.9%

Table 1: Results from the two independent analyses (first three rows) and from the combined fit of the two spectra (second six rows).




Figure 2: $\text{BR}(\phi \rightarrow K\bar{K}\gamma)$: excluded region (dark band) at 90% CL compared with theoretical estimates and the KLOE prediction (light band).

search for $[a_0(980) + f_0(980)] \rightarrow K\bar{K}$. This process has never been observed. Detection of a $K_S K_S$ pair is a clear signature of a $K\bar{K}$ pair with quantum numbers $J^{PC} = 0^{++}$, thus the $\phi \rightarrow K_S K_S \gamma \rightarrow 2(\pi^+\pi^-)\gamma$ process¹¹ is searched for, where the main background are the resonant $e^+e^- \rightarrow \phi\gamma \rightarrow K_S K_L \gamma$ and the continuum $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$ processes. From an integrated luminosity of 2.2 fb⁻¹, 5 candidate events are found in data, whereas 3 events are expected from Monte Carlo background samples. This leads to $\text{BR}(\phi \rightarrow K\bar{K}\gamma) < 1.9 \times 10^{-8}$ at the 90% CL. Fig. 2 shows the predictions of various theoretical models for the branching ratio. Some of them are excluded by this measurement. The present upper limit is consistent with the $\text{BR}(\phi \rightarrow K\bar{K}\gamma)$ prediction computed with $a_0(980)$ ⁸, $f_0(980)$ ^{12,13} couplings measured by KLOE.

search for $\gamma\gamma \rightarrow \sigma(600) \rightarrow \pi^0\pi^0$. While there is a long debate on the observation^{14,15} of the $\sigma(600)$ as a bound $\pi^+\pi^-$ state, there is no direct evidence for the $\sigma(600) \rightarrow \pi^0\pi^0$ decay. At DAΦNE, the detection of the process $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$ implies¹⁶ the intermediate process $\gamma\gamma$ to a scalar meson state. From a sample of 11 pb⁻¹ of data taken at $\sqrt{s} = 1$ GeV, the feasibility study of the $\pi^0\pi^0 \rightarrow 4\gamma$ invariant mass ($M_{4\gamma}$) spectrum – without tagging e^+ or e^- in the final state – is performed. Preliminary results show an excess of events in the $M_{4\gamma}$ region below 400 MeV, difficult to be explained by ϕ decays or e^+e^- annihilation processes. This preliminary work is encouraging and motivates the analysis extension to the whole sample of 240 pb⁻¹ collected by KLOE at $\sqrt{s} = 1$ GeV.

4 Conclusions

The KLOE data set, together with a precise simulation of the detector response and of a large number of several processes – ϕ decays and continuum e^+e^- annihilations – allowed for an extensive study of properties of scalar and pseudoscalar mesons:

- the most precise $\text{BR}(\eta \rightarrow \pi^+\pi^-e^+e^-)$ determination and the first measurement of the asymmetry between $\pi^+\pi^-$ and e^+e^- planes, found compatible with zero;
- evidence of valence gluons in the η' wavefunction to three standard deviations;
- good agreement between two $\eta\pi^0\gamma$ analyses with different systematics, where the combined fit of the two spectra points to a sizeable strange quark content in the $a_0(980)$;
- upper limit of the $\text{BR}(\phi \rightarrow K\bar{K}\gamma)$ at 90% CL with the whole KLOE statistics;
- excess of events with respect to known backgrounds in the 4γ invariant mass.

The programme of the KLOE-2 experiment, expected to roll in the e^+e^- interaction region of DAΦNE in the second half of 2009, includes meson spectroscopy with $\gamma\gamma$ interactions, improved measurements in the η - η' sector and observation of the $K\bar{K}\gamma$ final states.

References

1. M. Adinolfi *et al.* [KLOE Collaboration], Nucl. Instr. Meth. A **488**, 51 (2002).
2. M. Adinolfi *et al.* [KLOE Collaboration], Nucl. Instr. Meth. A **482**, 364 (2002).
3. C. Q. Geng, J. N. Ng and T. H. Wu, Mod. Phys. Lett. A **17**, 1489 (2002).
4. D. N. Gao, Mod. Phys. Lett. A **17**, 1583 (2002).
5. F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **675**, 283 (2009).
6. F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **648**, 267 (2007).
7. F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **669**, 223 (2008).
8. F. Ambrosino *et al.* [KLOE Collaboration], arXiv:0904.2539 [hep-ex].
9. N. N. Achasov and V. V. Gubin, Phys. Rev. D **56**, 4084 (1997).
10. G. Isidori, L. Maiani, M. Nicolaci and S. Pacetti, JHEP **0605**, 049 (2006).
11. F. Ambrosino *et al.* [KLOE Collaboration], arXiv:0903.4115 [hep-ex], submitted to PLB.
12. F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **634**, 148 (2006).
13. F. Ambrosino *et al.* [KLOE Collaboration], Eur. Phys. J. C **49**, 473 (2007).
14. E. M. Aitala *et al.* [E791 Collaboration], Phys. Rev. Lett. **86**, 770 (2001).
15. M. Ablikim *et al.* [BES Collaboration], Phys. Lett. B **598**, 149 (2004).
16. F. Nguyen, F. Piccinini and A. D. Polosa, Eur. Phys. J. C **47**, 65 (2006).